

**COMPLEXITY OF FREE RADICAL METABOLISM IN HUMAN ERYTHROCYTES**

## KOMPLEKSNOST METABOLIZMA SLOBODNIH RADIKALA U HUMANIM ERITROCITIMA

Aleksandra Nikolić-Kokić, Duško Blagojević, Mihajlo B. Spasić

University of Belgrade, Institute for Biological Research »Siniša Stanković«, Department of Physiology

**Summary:** The auto-oxidation of oxyhaemoglobin to methaemoglobin generating superoxide anion radical ( $O_2^{\cdot-}$ ) represents the main source of free radicals in the erythrocytes. Hydrogen peroxide is produced by  $O_2^{\cdot-}$  dismutation or originates from the circulation. Human erythrocytes are also exposed to the prooxidative actions of nitric oxide (NO) from circulation. Free radicals that may induce reactions with direct dangerous consequences to erythrocytes are also preceded by the reaction of  $O_2^{\cdot-}$  and NO producing peroxynitrite. In physiological settings, erythrocytes show a self-sustaining activity of antioxidative defence (AD) enzymes, such as: superoxide dismutase (SOD, EC 1.11.1.6), catalase (CAT, EC 1.11.1.6), glutathione peroxidase (GSHPx, EC 1.11.1.9) and glutathione reductase (GR, EC 1.6.4.2), as well as low molecular weight antioxidants: glutathione and vitamins E and C. Their coordinate actions protect the erythrocyte's biomacromolecules from free radical-mediated damage. Since there is no *de novo* synthesis of AD enzymes in mature erythrocytes, their defence capacity is limited. Free radicals influence antioxidative enzymes capacities and relative share of particular components in the whole antioxidative system. Therefore, by measuring changes in the activity of individual AD components, as well as their interrelations by statistical canonical discriminant methods, valuable data about the complexity, overall relations and coordinated actions in the AD system in erythrocytes and its relevance for systemic effects can be acquired.

**Keywords:** free radicals, superoxide anion radical, nitric oxide, antioxidants

**Kratak sadržaj:** Produkcija slobodnih radikala u eritrocitima uglavnom se odnosi na nastajanje superoksid anjon radikala ( $O_2^{\cdot-}$ ) putem autooksidacije oksihemoglobina u methemoglobin. Ljudski eritrociti izloženi su prooksidacionom delovanju vodonik-peroksida nastalog dismutacijom  $O_2^{\cdot-}$  ili iz cirkulacije, kao i azot oksidu (NO) iz cirkulacije. Od direktnih reakcija slobodnih radikala, reakcija  $O_2^{\cdot-}$  i NO uz nastajanje peroksinitrita je reakcija sa primarno štetnim posledicama po eritrocite. U eritrocitima se nalaze enzimi zaštite od oksidacionih oštećenja, kao što su superoksid dismutaza (SOD, EC 1.15.1.1), katalaza (CAT, EC 1.11.1.6), glutation peroksidaza (GSHPx, EC 1.11.1.9) i glutation reduktaza (GR, EC 1.6.4.2) kao i komponente male molekulske mase (glutation, vitamini E i C). Njihovim sadejstvom se kanališu reakcije slobodnih radikala tako da direktna oštećenja biomakromolekula budu što manja. Međutim, kako nema *de novo* sinteze enzima u maturiranim eritrocitima, kapacitet ovih sistema je ograničen, jer slobodnoradikalске vrste i direktno inhibiraju neke od enzima. Promene na enzimima i njihova inhibicija slobodnim radikalima utiču na kapacitet zaštite od oksidacionih oštećenja i relativni udeo pojedinih komponenti u ukupnom antioksidativnom potencijalu. To se može pratiti i preko promena aktivnosti pojedinačnih komponenti, ali i međusobnih odnosa između komponenti antioksidativne odbrane diskriminacionim statističkim metodama, koje ukazuju na sveukupnost i kompleksnost odnosa antioksidativnih komponenti u eritrocitima i njihov sistemski značaj.

**Ključne reči:** slobodni radikali, superoksid, azot-oksidi, antioksidanti

**Introduction**

The concept of an antioxidant defence system (ADS), as a means of preventing oxidative cell damage and promoting erythrocyte antioxidative instead of prooxidative role in the circulation, implies balanced activities of the erythrocyte ADS. Erythrocytes are exposed to oxygen radicals that are continuously generated primarily due to the auto-oxidation of oxyhaemoglobin (Hb) to methaemoglobin (1, 2). Erythrocytes

Address for correspondence:

Mihajlo B. Spasić  
Department of Physiology  
Institute for Biological Research »Siniša Stanković«  
Bulevar Despota Stefana 142, 11000 Belgrade, Serbia  
Tel: + 381 11 2078396; Fax: +381 11 2764422  
e-mail: spasa@ibiss.bg.ac.rs

are also exposed to oxidative pressure from plasma (particularly hydrogen peroxide –  $\text{H}_2\text{O}_2$  and nitric oxide –  $\text{NO}$ ). Under normal conditions erythrocytes contain sufficient levels of scavenger enzymes such as Cu,Zn-SOD, CAT, and selenium-dependent GSH-Px to protect themselves from free radical injury. Cu,Zn-SOD catalyses the dismutation of superoxide ( $\text{O}_2^{\cdot-}$ ) to  $\text{H}_2\text{O}_2$ , which is then independently converted to water by CAT or by GSH-Px (3). It is important to note that the activities of these enzymes in erythrocytes are higher than in most other tissues in the body (4). In humans, erythrocytes lack a nucleus, mitochondria and other organelles. Hence, there is no *de novo* synthesis of ADS components, making it difficult to maintain erythrocyte membranes intact for a long period – 120 days. A comprehensive understanding of the ADS should be based on the knowledge of activities and mutual interactions of the enzymes involved in free radical detoxication (5). Balanced action of antioxidant components is necessary for ROS homeostasis and appropriate redox state. Furthermore, if there is any coordinated action between the components, there might be a statistically significant correlation (positive and negative) between them. Therefore, changes in the activity of some antioxidant component should be accompanied by changes in other antioxidative enzymes in a correlated manner. Since correlation analysis calculates the relations between individual components, one can perform an alternative statistical test: canonical discriminant analysis, which calculates differences between groups taking into consideration the complete correlation matrix (i.e. composition of antioxidant defence) of separate groups. The ADS in human erythrocytes with preserved homeostasis finely retunes its composition according to plasma oxidative demands. An increase in the level of a specific plasma lipid component may potentiate membrane lipid peroxidation in erythrocytes and decrease intra-erythrocyte production of  $\text{O}_2^{\cdot-}$ , which could result in a negative correlation between SOD and GSH-Px activities found in some experiments (6–9). The discovery of the haemoglobin-cholesterol (Hb-Ch) complex implies the way in which cholesterol may influence the organisation of ADS in erythrocytes (7).

Physiologically active molecules (such as nitric oxide ( $\text{NO}$ )) can also react with ROS. The reaction between  $\text{O}_2^{\cdot-}$  and  $\text{NO}$  naturally occurs in cells producing peroxynitrite ( $\text{ONOO}^-$ ) which is considered to be a cytotoxic molecule. The production of so-called reactive nitrogen species (RNS) represents a consequence of  $\text{ONOO}^-$  formation. RNS provokes nitrosative modifications of molecules and can induce nitrosative stress (10). The interactions of ROS and RNS with antioxidative enzymes leading to the inhibition of their activities, represent the primary cause of disturbed correlations between their activities in erythrocytes. If ROS production overwhelms the ADS, oxidative stress occurs leading to oxidative damages. Non-specific interactions of ROS with different classes of intra-

cellular molecules induce oxidative damage leading to the impairment of cellular homeostasis. As accurate detection of *in vivo* ROS concentrations is still problematic, changes in the ADS may serve as a good indicator of processes within the organism, as it responds to ROS production, changes of the environment (as low environmental temperature) (11) and pathological conditions (9). It should be stressed that the ADS is species- and tissue-specific (12) and that antioxidative enzymes in erythrocytes and the dominant source of ROS in erythrocytes are in many aspects quite different in comparison with extracellular fluids and other tissues.

#### *Antioxidative enzymes in erythrocytes*

Human erythrocytes are the most abundant and one of the most specialized cells in the body. The main function of erythrocytes is the transport of oxygen ( $\text{O}_2$ ) and mediation of carbon dioxide ( $\text{CO}_2$ ) production (13). As the red blood cell emerges from the bone marrow, it loses its nucleus, ribosomes, and mitochondria, and therefore the capacity for cell division, protein synthesis, and mitochondrial-based oxidative reactions (13, 14). As a consequence of their physiologic role, erythrocytes are exposed to continuous oxidative stress. Although oxidative stress may damage the red cell itself, the mass effect of large quantities of ROS leaving the red cell could have a tremendous potential to damage other components of the circulation (15), and *vice versa*.

Normal erythrocytes have been shown to have a reducing capacity that is 250 times greater than their oxidizing potential (16). However, in some pathological conditions, the erythrocyte ADS seems to be insufficient (17, 18). Erythrocytes contain a significant quantity of CuZn SOD, which keeps intra-erythrocyte  $\text{O}_2^{\cdot-}$  levels at concentrations as low as  $10^{-13}$  mol/L. The GSH dependent enzymes (GSH-Px and GR) are also normally present, enabling removal of  $\text{H}_2\text{O}_2$  by oxidation/reduction of GSH. No manganese superoxide dismutase (MnSOD) is present, as there are no mitochondria in erythrocytes.

Erythrocytes can act as sinks for extracellular  $\text{H}_2\text{O}_2$  (freely crosses membranes) and  $\text{O}_2^{\cdot-}$  as it can enter the cells via anion channels. These channels have different functions: anchoring the cytoskeleton and exchanging  $\text{HCO}_3^-$  for  $\text{Cl}^-$ . However,  $\text{O}_2^{\cdot-}$  and  $\text{NO}_2^-$ , as well as the very toxic  $\text{ONOO}^-$  can also enter erythrocytes through these channels. Once inside the cell,  $\text{ONOO}^-$  reacts with GSH and oxyhaemoglobin, yielding  $\text{NO}_3^-$ , methaemoglobin, some  $\text{H}_2\text{O}_2$  (via  $\text{O}_2^{\cdot-}$  dismutation), and some ferrylhaemoglobin.

Unfortunately, erythrocytes are particularly vulnerable to oxidative stress due to constant exposure to endogenously (autooxidation of haemoglobine-Hb) and exogenously generated radicals ( $\text{H}_2\text{O}_2$  and  $\text{NO}$  from plasma). As well as other erythrocyte components,


antioxidative enzymes are exposed to the prooxidative pressure of  $O_2^{\cdot-}$ ,  $H_2O_2$ ,  $ONOO^-$ . In the reaction with different radicals, antioxidant enzymes may lose their primary activity and gain prooxidative properties, which can lead to enhanced oxidative pressure and generalised oxidative stress in the whole body.

#### *Inhibition of antioxidant enzymes*

The examination of AD enzyme activities *in vitro* has shown that CuZn-SOD possesses a constant specific activity (activity per mg of purified enzyme) and may be inhibited irreversibly by CN<sup>-</sup> or reversibly by  $H_2O_2$  or by copper chelators such as DDC (diethyldithiocarbamate) (19). Multiple electroforetic profiles of CuZn-SOD develop as a consequence of the enzyme aging (20). Hydrogen peroxide, or rather its conjugate base ( $OH_2^-$ ), reacts with CuZnSOD, reducing Cu(II) to Cu(I), followed by the reaction of Cu(I) with a second hydrogen peroxide. This leads to the oxidation of the active site, putatively described as copper-bound hydroxyl radical (21). This, in turn, leads to the inactivation of enzyme through 2-oxohistidine formation (22) and to the oxidation of various substrates in the enzyme's behaviour known as the »peroxidative« activity of CuZn-SOD (21–24). DDC is known to be a potent Cu chelating agent and is one of the most widely used SOD inhibitors both *in vivo* and *in vitro* (25–28). The mechanism of DDC-mediated CuZn-SOD inhibition has been described (29). DDC-mediated SOD1 inhibition could serve as a screen for oxidatively damaged SOD 1 protein in the blood of acute myocardial infarction subjects. Oxidative inactivation of red cell SOD by its product  $H_2O_2$  generates a modified protein which is recognised and selectively degraded by an intracellular proteolytic pathway (30). Both the loss of SOD activity and modified binding of copper by the active site appear to precede proteolytic recognition and degradation. The selective degradation of  $H_2O_2$ -modified SOD in red cell extracts is now seen to be catalysed by an ATP-independent proteolytic pathway. Oxidised histidine 118 to 2-oxo histidine is generated at the active site of CuZn-SOD exposed to  $H_2O_2$  (22). An increased concentration of  $H_2O_2$  allows the production of hydroxyl radicals, especially in the presence of catalytically-active metals (23, 31).

Oxidatively damaged SOD may cause further increase in free radicals due to more solvent exposed  $Cu^+$ . Such a situation is presented in *Figure 1*. Electron paramagnetic resonance spectra of SOD inhibited with  $H_2O_2$  show that 60 minutes after the application of  $H_2O_2$ , oxidatively damaged SOD catalyse the Fenton reaction and promote the generation of hydroxyl radicals.

Similarly to CuZn-SOD, other antioxidative enzymes can also be changed or inactivated by reactive oxygen species. Pigeolet et al. (32) tested the effect of



**Figure 1** Electron paramagnetic resonance spectra of hydroxyl radicals 'trapped' with a DEPMPO spin-trap in the system composed of CuZn-SOD and hydrogen peroxide (5 mmol/L). In the min lane it is evident that an intensive Fenton reaction occurs.

$H_2O_2$ , cumene hydroperoxide, t-butyl hydroperoxide and  $OH^{\cdot}$  and  $O_2^{\cdot-}$  on GSH-Px, CuZn SOD and CAT. The activity of GSH-Px was decreased by 50% inactivation in the presence of hydrogen peroxide (0.1 M), cumene hydroperoxide ( $3 \times 10^{-4}$  mmol/L), and t-butyl hydroperoxide ( $5 \times 10^{-5}$  mol/L). Unlike  $OH^{\cdot}$ ,  $O_2^{\cdot-}$  did not inactivate this enzyme. CAT was inactivated by NO,  $OH^{\cdot}$  and by  $O_2^{\cdot-}$  but organic peroxides had no effect. Similar to our results, Pigeolet showed a 50% SOD inhibition with  $4 \times 10^{-4}$  mmol/L  $H_2O_2$  (32).

Clearly, the ADS is affected by an imbalance between the production of reactive oxygen species and their decomposition, which leads to new ADS settings. Our scientific research has been directed towards finding a cross-relation between the ADS profile of easily accessible human material such as erythrocytes, and the prooxidative changes at the primary site of radical generation in other tissues. In other words, how a changed ADS composition implicates the redox processes in pathological conditions. To answer the above questions, we applied the mathematical model of canonical discriminated analysis to ADS changes measured in the erythrocytes of patients with three different diseases: cardiovascular (AIM), neurological (ALS) and psychiatric (SCH).

#### *Acute myocardial infarction (AMI)*

A disturbed balance between the production of both reactive oxygen and nitrogen species and their elimination as a consequence of acute myocardial infarction (AMI) has been postulated to represent the molecular basis of oxidative stress and damage which are important factors in reperfusion injury (33). During myocardial reperfusion injury the production of reactive oxygen species arises via several routes including mitochondrial respiratory-chain enzymes, xanthine oxidase, and non-phagocytic and neutrophil NADPH oxidase (34). An additional source of reactive oxygen species during heart failure may be angiotensin II and catecholamines (more precisely, the auto-oxidation of catecholamines), whose increased levels represent a consequence of sympathetic activation. An excess of  $H_2O_2$  and NO in extracellular medium is followed by

the rise of intracellular level due to the fact that these species can permeate through biological membranes.  $H_2O_2$  and NO may affect the ADS by inhibiting CuZn-SOD (20) and CAT (35). This results in conditions that propagate reperfusion injury via oxidative changes in erythrocytes related to the shift of enzyme behavior from antioxidant to prooxidant. Tsao and colleagues have shown that endothelial cells lose their function 2.5 min after reperfusion injury by a mechanism implicating reactive oxygen species formation (36). Mitochondria in an ischemic heart may additionally increase the generation of hydrogen peroxide during reperfusion (37). Hydrogen peroxide may diffuse from myocytes into the bloodstream to affect erythrocyte function.

Presented settings imply strong correlation between the ADS of erythrocytes and oxidative conditions in the heart and vessels. However, in some pathophysiologies this may not be the case. During reperfusion induced by streptokinase treatment, no significant correlations between the antioxidant defence enzyme activities were apparent (38), while in the control population we found a significant positive correlation between the activities of CAT and GSH-Px in erythrocytes (38, 39). This indicated that ischaemia/reperfusion had disturbed the coordinated action of AD enzymes in the erythrocytes of ALM patients. Only after sustained carvedilol therapy (168 hours duration) a significant correlation between CuZnSOD and CAT was found, indicating that carvedilol therapy had a positive effect on re-establishing the normal relationship between antioxidant setup in the erythrocytes and heart. This is supported by the results of the two-way ANOVA that showed a significant effect of treatment (38).


*Amyotrophic lateral sclerosis (ALS) –  
a fatal progressive disorder*

Amyotrophic lateral sclerosis (ALS), often called motor neuron disease (MND), is an adult-onset neurodegenerative disease characterised by progressive injury and death of lower motor neurons in the spinal cord and brainstem, and upper motor neurons in the motor cortex (40). The primary cause of disease is unknown, and the mechanism of motor neuron injury is complex and incompletely understood. In 1993 Rosen and co-workers found that 20% of familial MND cases (5–10% of all cases are familial) are caused by mutations in copper/zinc superoxide dismutase (41). Since that time several hypothetical mechanisms have been predicted: oxidative stress, excitotoxicity caused by aberrant glutamate signaling, mitochondrial dysfunction, disruption of the neurofilament network and intracellular trafficking along neurofilament aggregation of proteins, and the involvement of non-neuronal cells in the vicinity of motor neurons. Events in these mechanisms culminate in a caspase-dependent programmed cell death pathway resembling apoptosis (42). Recent findings corroborated the

hypothesis that glutamate-mediated NO overproduction plays an important role in the pathogenesis of ALS (43). Due to their ability to permeate through biological membranes, an excess of NO and  $H_2O_2$  may be present in the surrounding media of motor neurones (44). Some data suggest that the ADS activity is altered in erythrocytes from ALS patients (45, 46) before and after adjuvant ALS therapy (47). Antioxidative defense enzymes in erythrocytes are capable of detoxifying reactive oxygen species (produced endogenously or exogenously), but as mentioned in the preceding sections, the enzymes may be structurally modified and inactivated by reactive oxygen and nitrogen species. Both balanced and coordinated ADS activities are of the utmost importance for their correct physiological function.

In the erythrocytes of SALS patients a significant negative correlation between GSH-Px and CAT was found (39). However, in FALS patients (with mutated SOD) and healthy subjects with mutations in the SOD molecule, no significant correlations were found (unpublished data). These mean that the basic relationships between the ADS components in ALS are different and depend on systemic conditions, i.e. the amount of produced ROS and RNS.

Using the canonical discriminant analysis, it is possible to discriminate different categories (in this study, the analysed groups) according to the composition of the antioxidative defense components, and to determine which component significantly contributes to this difference (48, 49). In this way the antioxidative erythrocyte enzymes in SALS, FALS patients, asymptomatic carriers and controls were analyzed by two-way ANOVA and canonical discriminant analyses. The results obtained showed that all examined AOS enzymes significantly contributed to the difference in antioxidative defense composition observed between the groups (Figure 2).




**Figure 2** Two-dimensional discriminant analysis showing differences in the ADS activities in erythrocytes from SALS (–/+), from FALS (+/+), from asymptomatic carriers of SOD1 (Leu144Phe mutation) (+/-) and from the control group (–/-).

### Schizophrenia

Recent findings suggest that multiple neurotransmitter systems may be faulty in schizophrenic patients (50). Metabolic products of such a non-functional system, such as  $H_2O_2$  and RNS, may inhibit erythrocyte AD enzymes. From the initial studies of antioxidant defence in schizophrenia (Sch.) (51) to the most recent one (52), disturbed balance in the activity of antioxidant defence enzymes in the erythrocytes of schizophrenics was found. In never-medicated first-episode psychotic patients, lower levels of SOD activity were reported, but no change was observed in the activities of CAT and GSH-Px (53). According to our findings, it appears that schizophrenia creates conditions that increase the level of  $H_2O_2$  affecting circulating cells.

Our idea in examining the ADS erythrocyte enzymes in Sch patients was to investigate whether systemic oxidative misbalance reflects on the composition of the ADS of erythrocytes from Sch type I (Sch I) and type II (Sch II) patients. The correlation analysis of antioxidant defence enzymes showed a significant negative correlation between GSH-Px and CAT activities in patients with Sch I. In patients with Sch II, GSH-Px activity showed a statistically significant positive correlation with GR. Canonical discriminant analysis distinguished Sch I and Sch II patients from the controls, and among each other, with a high degree of certainty, according to the overall group composition of AD enzymes (Figure 3) (54). The results indicated a difference in the composition of ADS between controls and antipsychotic treated Sch I and Sch II patients, with a possible negative feedback influence on the pathological process, and may represent a rationale for applying antioxidants in Sch therapy.

According to our findings, it appears that schizophrenia creates conditions that increase the level of hydrogen peroxide, affecting circulating cells. The source of  $H_2O_2$  in the circulation may be an increased monoamine oxidase activity, since an increased turnover rate of catecholamines was found in schizophrenic patients. Such changes in the circulation may shift erythrocyte role from antioxidative to prooxidative. Presumably the best way to address these negative effects is through controlling oxidative stress in a physiological manner, not aiming only at the level of antioxidants, but rather at their interactions. According to Crow (55), Sch I and Sch II show different pathogenetic mechanisms which result in differences in the therapeutic response to antipsychotics. Crow wrote that there were two syndromes, due to different pathophysiological processes: Sch I with changes in dopamine transmission and Sch II with encephalopathy (55). Our



**Figure 3** Two-dimensional discriminant analysis showing differences in the activities of antioxidant defense enzymes in the erythrocytes of controls, Sch I and Sch II.

results on ADS changes seem to confirm such a statement.

### Conclusion

We first used antioxidative defence enzyme relations as a possible bioindicator of the effects of ionizing radiation (56). Since then the concept that the oxidative status of various tissues may be determined by studying the antioxidant defence system has been verified in several papers. Erythrocytes are particularly vulnerable to oxidative stress because they are exposed to oxygen radicals, that are continuously generated primarily due to the auto-oxidation of haemoglobin (Hb). There is a defence system against oxidative stress in erythrocytes composed of CuZn-SOD, CAT, selenium-dependent GSH-Px, and GR. Erythrocytes are also exposed to oxidative pressure from plasma (particularly mediated by  $H_2O_2$  and NO). The measurements of changes in the activities of individual antioxidative defence components as well as their interrelations, using statistical canonical discriminant methods, are capable of providing a valuable insight into the complexity of overall relations, coordinated actions in the antioxidative defence system in erythrocytes and its relevance to systemic effects.

### Conflict of interest statement

The authors stated that there are no conflicts of interest regarding the publication of this article.

## References

- Rifkind JM, Nagababu E, Ramasamy S, Ravi LB. Hemoglobin redox reactions and oxidative stress. *Redox Rep* 2003; 8 (5): 234–7.
- Winterbourn CC. Haemoglobin oxidation and free radical production in the red cell. *Biomed Biochim Acta* 1983; 42 (11–12): S134–8.
- Chance B, Sies H, Boveris A. Hydroperoxide metabolism in mammalian organs. *Physiol Rev* 1979; 59 (3): 527–605.
- Spasić MB. Antioxidative defence in mammals – A review. *Jugoslav Med Biochem* 1993; 12: 1–9.
- Guemouri L, Artur Y, Herbeth B, Jeandel C, Cuny G, Siest G. Biological variability of superoxide dismutase, glutathione peroxidase, and catalase in blood. *Clin Chem* 1991; 37 (11): 1932–7.
- Nikolić M, Vranić D, Spirić A, Batas V, Nikolić-Kokić A, Radetić P, et al. Could cholesterol bound to haemoglobin be a missing link for the occasional inverse relationship between superoxide dismutase and glutathione peroxidase activities? *BBRC* 2006; 348: 265–70.
- Nikolić M, Nikolić-Kokić A, Stanić D, Blagojević DP, Vranić D, Jones D, Niketić V, Spasić MB. Does cholesterol bound to haemoglobin affects the antioxidative enzyme defence system in human erythrocytes? *J Serb Chem Soc* 2007; 72: (4): 339–45.
- Kasimanickam R, Kasimanickam V, Thatcher CD, Nebel RL, Cassell BG. Relations among lipid peroxidation, glutathione peroxidation, glutathione peroxidase, superoxide dismutase, sperm parameters, and competitive index in dairy bulls. *Theriogenology* 2007; 67 (5): 1004–12.
- Nikolić-Kokić A, Stević Z, Blagojević D, Davidović B, Jones DR, Spasić MB. Alterations in anti-oxidative defence enzymes in erythrocytes from sporadic amyotrophic lateral sclerosis (SALS) and familial ALS patients. *Clin Chem Lab Med* 2006; 44 (5): 589–93.
- Halliwell B, Gutteridge JMC. In: *Free Radicals in Biology and Medicine*, Fourth edition. New York, Oxford University Press Inc., 2007.
- Blagojević D. Antioxidant systems in supporting environmental and programmed adaptations to low temperatures. *Cryo Letters* 2007; 28 (3): 137–50.
- Blagojević D, Buzadžić B, Korać B, Saičić ZS, Radojičić R, Spasić MB, Petrović VM. Seasonal changes in the anti-oxidative defence in ground squirrels (*Citellus citellus*): possible role of GSH-Px. *J Environ Pathol Toxicol Oncol* 1998; 17 (3–4): 241–50.
- Spasojević I. Electron paramagnetic resonance – A powerful tool of medical biochemistry in discovering mechanisms of disease and treatment prospects. *Journal of Medical Biochemistry* 2010; 29: 175–188.
- Bunn HF. Pathophysiology of the anemias. In: Wilson JD, Braunwald E, Isselbacher KJ, editors. *Harrison's Principle of Internal Medicine*. New York: McGraw-Hill Inc; 1991; 1514–8.
- Johnson RM, Goyette Jr G, Ravindranath Y, Ho YS. Hemoglobin autoxidation and regulation of endogenous  $H_2O_2$  levels in erythrocytes. *Free Radic Biol Med* 2005; 39: 1407–17.
- Scott, MD, Eaton JW, Kuypers FA, Chiu DT, Lubin BH. Enhancement of erythrocyte superoxide dismutase activity: effects on cellular oxidant defense. *Blood* 1989; 74, 2542–9.
- Chakraborty D, Bhattacharyya M. Antioxidant defense status of red blood cells of patients with beta-thalassemia and Ebeta-thalassemia. *Clin Chim Acta* 2001; 305: 123–9.
- Tesoriere L, D'Arpa D, Butera, D, Allegra M, Renda D, Maggio A, Bongiorno A, Livrea MA. Oral supplements of vitamin E improve measures of oxidative stress in plasma and reduce oxidative damage to LDL and erythrocytes in beta-thalassemia intermedia patients. *Free Radic Res* 2001; 34: 529–40.
- Padmaja S, Squadrito GL, Pryor WA. Inactivation of glutathione peroxidase by peroxynitrite. *Arch Biochem Biophys* 1998; 349: 1–6.
- Mavelli I, Ciriolo MR, Rotilio G. Multiple electrophoretic variants of Cu, Zn superoxide dismutase as expression of the enzyme aging. Effects of  $H_2O_2$ , ascorbate and metal ions. *Biochem Biophys Res Commun* 1983; 117 (3): 677–81.
- Hodgson EK, Fridovich I. The interaction of bovine erythrocyte superoxide dismutase with hydrogen peroxide: chemiluminescence and peroxidation. *Biochemistry* 1975; 14 (24): 5299–303.
- Uchida K, Kawakishi S. Identification of oxidized histidine generated at the active site of Cu,Zn-superoxide dismutase exposed to  $H_2O_2$ . Selective generation of 2-oxo-histidine at the histidine 118. *J Biol Chem* 1994 269 (4): 2405–10.
- Yim MB, Chock PB, Stadtman ER. Copper, zinc superoxide dismutase catalyzes hydroxyl radical production from hydrogen peroxide. *Proc Natl Acad Sci USA* 1990; 87 (13): 5006–10.
- Jewett SL, Olmsted HK, Marach JA, Rojas F, Silva K. Anion protection of CuZnSOD during peroxidative activity with  $H(2)O(2)$ . *Biochem Biophys Res Commun* 2000; 274 (1): 57–60.
- Heikkila RE, Cabbat FS, Cohen G. In vivo inhibition of superoxide dismutase in mice by diethyldithiocarbamate. *J Biol Chem* 1976; 251 (7): 2182–5.
- Radojičić R, Borković S, Pavlović S, Nikolić A, Blagojević D, Saičić S, Spasić M. The effect of diethyldithiocarbamate on antioxidant enzyme activities in the blood of rats. *Acta Vet* 2002; 52: 329–36.
- Radojičić R, Spasić M, Milić B, Saičić ZS, Petrović VM. Age-related differences in the effect of diethyldithiocarbamate and cyclohexamide on the liver copper, zinc-containing superoxide dismutase in the rat. *Jugoslav Physiol Pharmacol Acta* 1987; 23: 227–33.
- Oreščanin-Dusić Z, Milovanović S, Blagojević D, Nikolić-Kokić A, Radojičić R, Spasojević I, Spasić M. Diethyldithiocarbamate potentiates the effects of protamine sulphate in the isolated rat uterus. *Redox Rep* 2009; 14 (2): 48–54.

29. Iciek M, Włodek L. Biosynthesis and biological properties of compounds containing highly reactive, reduced sulfane sulfur. *Pol J Pharmacol* 2001; 53 (3): 215–59.
30. Davis CA, Hearn AS, Fletcher B, Bickford J, Garcia JE, Leveque V, et al. Potent anti-tumor effects of an active site mutant of human manganese-superoxide dismutase. Evolutionary conservation of product inhibition. *J Biol Chem* 2004; 279 (13): 12769–76.
31. Peled-Kamar M, Lotem J, Wirguin I, Weiner L, Hermalin A, Groner Y. Oxidative stress mediated impairment of muscle function in transgenic mice with elevated level of wild-type Cu/Zn superoxide dismutase. *Proc Natl Acad Sci USA* 1997; 94 (8): 3883–7.
32. Pigeolet E, Corbisier P, Houbion A, Lambert D, Michiels C, Raes M, Zachary MD, Remacle J. Glutathione peroxidase, superoxide dismutase, and catalase inactivation by peroxides and oxygen derived free radicals. *Mech Ageing Dev* 1990; 51 (3): 283–97.
33. Simović M, Spasić MB, Michelson AM. Free radicals in human myocardial reperfusion injury. *Life Chem Rep* 1995; 12: 227–70.
34. Grieve DJ, Shah AM. Oxidative stress in heart failure. More than just damage. *Eur Heart J* 2003; 24 (24): 2161.
35. Brunelli L, Yermilov V, Beckman JS. Modulation of catalase peroxidatic and catalytic activity by nitric oxide. *Free Radic Biol Med* 2001; 30 (7): 709–14.
36. Tsao PS, Aoki N, Lefer DJ, Johnson G 3rd, Lefer AM. Time course of endothelial dysfunction and myocardial injury during myocardial ischemia and reperfusion in the cat. *Circulation* 1990; 82 (4): 1402–12.
37. Turrens JF, Beconi M, Barilla J, Chavez UB, McCord JM. Mitochondrial generation of oxygen radicals during reoxygenation of ischemic tissues. *Free Radic Res Commun* 1991; 12-13 Pt 2: 681–9.
38. Kastratović DA, Vasiljević ZM, Spasić BM, Peruničić PJ, Matić M, Blagojević PD, et al. Carvedilol Increases Copper-Zinc Superoxide Dismutase Activity in Patients with Acute Myocardial Infarction. *Basic & Clinical Pharmacology & Toxicology* 2007; 101: 138–42.
39. Nikolić A, Blagojević D, Stević Z, Niketić V, Saičić ZS, Spasić MB. Activities of AD enzymes in the blood of ALS patients-base for use of antioxidants in the treatment of ALS. *Proceeding of the 11 Biennial Meeting of the Society for Free Radical Research International 2002 by Monduzzi Editore S.p.A.–MEDIMOND Inc.* 323–6.
40. Barber SC, Mead RJ, Shaw PJ. Oxidative stress in ALS: a mechanism of neurodegeneration and a therapeutic target. *Biochim Biophys Acta* 2006; 1762 (11–12): 1051–67.
41. Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, et al. Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 1993; 362 (6415): 59–62.
42. Sathasivam S, Ince PG, Shaw PJ. Apoptosis in amyotrophic lateral sclerosis: a review of the evidence. *Neuropathol Appl Neurobiol* 2001; 27 (4): 257–74.
43. Tohgi H, Abe T, Yamazaki K, Murata T, Ishizaki E, Isobe C. Increase in oxidised NO products and reduction in oxidised glutathione in cerebrospinal fluid from patients with sporadic form of amyotrophic lateral sclerosis. *Neurosci Lett* 1999; 260 (3): 204–6.
44. Nikolić-Kokić A, Stević Z, Stojanović S, Blagojević PD, Jones DR, Pavlović S, Niketić V, Apostolski S, Spasić MB. Biotransformation of nitric oxide in the cerebrospinal fluid of amyotrophic lateral sclerosis patients. *Redox Report* 2005; 10 (5): 265–70.
45. Nikolić AL, Stević ZS, Blagojević D, Saičić ZS, Spasić MB. Activities of antioxidant defence enzymes in the blood of individuals with Leu144Phe mutation. *Jugoslav Med Biochem* 2005; 24 (2): 111–4.
46. Przedborski S, Donaldson DM, Murphy PL, Hirsch O, Lange D, Naini AB, et al. Blood superoxide dismutase, catalase and glutathione peroxidase activities in familial and sporadic amyotrophic lateral sclerosis. *Neurodegeneration* 1996; 5: 57–64.
47. Apostolski S, Marinković Z, Nikolić A, Blagojević D, Spasić MB, Michelson AM. Glutathione Peroxidase in Amyotrophic Lateral Sclerosis – the Effects of Selenium Supplementation. *J Environ Pathol Toxicol Oncol* 1998; 17 (2): 325–9.
48. Hinkle ED, Wiersma W, Jurs GS. *Applied statistics for behavioral sciences*, 3rd ed. Boston, MA: Houghton Mifflin Company, 1994.
49. Manley BFJ. *Multivariate statistical methods. A primer.* London: Chapman and Hall, 1986.
50. Du Bois TM, Deng C, Huang XF. Membrane phospholipid composition, alterations in neurotransmitter systems and schizophrenia. *Progress in Neuro Psychopharmacology Biological Psychiatry* 2005; 29: 878–88.
51. Michelson, AM. Medical aspects of superoxide dismutase. *Life Chemistry Reports* 1987; 6: 1–142.
52. Ranjekar PK, Hinge A, Hegde MV, Ghate M, Kale A, Sitasawad S, Wagh UV, Debsikdar VB, Mahadik SP. Decreased antioxidant enzymes and membrane essential polyunsaturated fatty acids in schizophrenic and bipolar mood disorder patients. *Psychiatry Res* 2003; 121 (2): 109–22.
53. Mukherjee S, Mahadik SP, Scheffer R, Correnti EE, Kelkar H. Impaired antioxidant defence at the onset of psychosis. *Schizophrenia Research* 1996; 19: 19–26.
54. Miljević C, Nikolić-Kokić A, Saičić ZS, Milosavljević M, Blagojević D, Lečić-Toševski D, et al. Correlation analysis confirms differences in antioxidant defence in the blood of type I and II schizophrenic male patients treated with anti-psychotic medication. *Psychiatry Research* 2010; 178: 68–72.
55. Crow TJ. Positive and negative schizophrenic symptoms and the role of dopamine. *Br J Psychiatry* 1980; 137: 385–6.
56. Žunić Z, Blagojević D, Spasić M, Đuić J, Nikolić A, Marković S, Saičić ZS. Low dose radiation effects on the activity of antioxidative defence in the blood of healthy examinees. *Health effects of low dose radiation. Challenges of the 21st century*, BNES, London 1997, 214–1.

Received: April 28, 2010

Accepted: May 26, 2010