

**ASSOCIATION OF HYPONATREMIA
AND HYPOVITAMINOSIS D IN AMBULATORY ADULTS**VEZA IZMEĐU HIPONATRIJEMIJE I HIPOVITAMINOZE D KOD
POKRETNIH BOLESNIKAGianfranco Cervellin¹, Gianluca Salvagno², Laura Bonfanti¹,
Patrizia Bonelli³, Gian Cesare Guidi², Giuseppe Lippi³¹Emergency Department, Academic Hospital of Parma, Parma, Italy²Laboratory of Clinical Chemistry and Hematology, University of Verona, Verona, Italy³Laboratory of Clinical Chemistry and Hematology, Academic Hospital of Parma, Parma, Italy**Summary**

Background: Hyponatremia and vitamin D deficiency are frequent disorders, and both have been associated with gait disturbances, falls and fractures. The aim of this study was to evaluate the existence of an association between serum sodium and vitamin D serum levels.

Methods: We performed a retrospective investigation to establish whether hyponatremia and vitamin D deficiency may be associated in a general population of unselected outpatients. An electronic search was performed in the laboratory information systems of the Hospital of Verona and the Hospital of Parma (Italy), to retrieve combined results for total vitamin D and sodium obtained in all outpatients referred for health check-up in the year 2013.

Results: Combined results of vitamin D and sodium could be retrieved for 5097 outpatients (3859 females and 1238 males; mean age 64±17 years). Vitamin D deficient subjects displayed significantly lower levels of serum sodium (140 versus 141 mmol/L; $p<0.001$), along with a significantly higher rate of hyponatremia (6.3% versus 5.1%; $p=0.037$). Accordingly, hyponatremic subjects had significantly lower levels of serum vitamin D (55 versus 60 nmol/L; $p=0.015$), along with a significantly higher rate of vitamin D deficiency (41.8% versus 36.1%; $p=0.030$). A highly significant correlation was found between sodium and total vitamin D after adjustment for age and gender ($p<0.001$).

Kratak sadržaj

Uvod: Hiponatrijemija i nedostatak vitamina D su česti poremećaji i povezani su sa teškoćama u hodu, padovima i prelomima. Cilj ove studije bio je da se utvrdi da li postoji povezanost između natrijuma u serumu i nivoa vitamina D u serumu.

Metode: Sproveli smo retrospektivno istraživanje da bismo utvrdili postojanje veze između hiponatrijemije i nedostatka vitamina D u opštoj populaciji bolesnika. Izvršena je elektronska pretraga laboratorijskih informacionih sistema Bolnice u Veroni i Bolnice u Parmi (Italija), kako bi se dobili kombinovani rezultati za ukupni vitamin D i natrijum za sve bolesnike koji su upućeni na zdravstveni pregled tokom 2013.

Rezultati: Kombinovani rezultati za vitamin D i natrijum dobijeni su za 5097 bolesnika (3859 žena i 1238 muškaraca; prosečna starost 64±17 godina). Kod pacijenata s nedostatkom vitamina D nađeni su značajno niži nivoi natrijuma u serumu (140 prema 141 mmol/L; $p<0,001$), kao i značajno viša stopa hiponatrijemije (6,3% prema 5,1%; $p=0,037$). U skladu s tim, bolesnici sa hiponatrijemijom imali su značajno niže nivoje vitamina D u serumu (55 prema 60 nmol/L; $p=0,015$), kao i značajno višu stopu deficijencije vitamina D (41,8% prema 36,1%; $p=0,030$). Pronađena je veoma značajna korelacija između natrijuma i ukupnog vitamina D posle prilagođavanja za uzrast i pol ($p<0,001$).

Zaključak: Rezultati ove studije prvi put su pokazali da postoji značajna korelacija između nivoa natrijuma u serumu i

Address for correspondence:

Dr. Gianfranco Cervellin
U.O. Pronto Soccorso e Medicina d'Urgenza
Azienda Ospedaliero-Universitaria di Parma,
Via Gramsci, 14
43126 - Parma, Italy
Tel. 0039-0521-703800
e-mail: gcervellin@ao.pr.it

Conclusions: The results of this study demonstrate for the first time the existence of a significant correlation between the serum levels of sodium and total vitamin D in a general population of unselected outpatients.

Keywords: hyponatremia, osteoporosis, sodium, vitamin D, elderly

Introduction

Both hyponatremia and vitamin D deficiency are frequent disorders in the elderly (1, 2). The former condition affect up to 15–30% of hospitalized patients (3), and approximately 50% of these cases are due to the syndrome of inappropriate antidiuretic hormone secretion (SIADH), whereas the remaining cases are mainly attributable to hyatrogenic effects (most frequently to diuretics, psychotropic drugs, anticonvulsants), or chronic conditions such as hypothyroidism, congestive heart failure, liver cirrhosis and renal failure (4). Hyponatremia, generally defined as a serum sodium concentration < 136 mmol/L (5) has been recently associated with gait disturbances, falls and fractures in the elderly (6). Recent data also show that hyponatremia may directly influence bone metabolism and can hence represent an important predisposing factor for osteoporosis and skeletal frailty (7). Therefore, this electrolyte disorder should be regarded as a direct contributor of osteoporosis and also as an important risk factor for impaired gait, finally leading to falls. In animal studies, sustained hyponatremia was effective to induce progressive reduction of bone mineral density and decreased bone ash calcium, along with reduced phosphate and sodium contents at the tibia and lumbar vertebrae. In these animals, the administration of vitamin D compensated for the reduction in bone formation and halted bone loss (8). The hyponatremic animals also developed hypogonadism, and markedly decreased testicular weight with abnormal testicular histology, along with decreased body fat, skeletal muscle sarcopenia, and cardiomyopathy (8). Interestingly, in a similar experiment, the administration of activated vitamin D was effective to mitigate the development of left ventricular alterations in hypertensive rats (9).

Interestingly, vitamin D deficiency seems to show similar features. Although there is no universal consensus on optimal levels of 25-hydroxyvitamin D (25-OHD, i.e., the main circulating form of the vitamin) as measured in serum, vitamin D deficiency is generally defined as a 25-OHD level of less than 20 ng/mL (50 nmol/L), whereas a level of 21 to 29 ng/mL (52 to 72 nmol/L) indicates a relative insufficiency, and a level of 30 ng/mL (75 nmol/L) or greater can be considered as normal (10). Vitamin D insufficiency and deficiency are highly prevalent, and this is mirrored by the fact that more than half of the worldwide population has levels of 25-OHD below 30 ng/mL (11).

Vitamin D is classically known for its role in bone health, playing an essential role in the maintenance

ukupnog vitamina D u opštoj populaciji neselektovanih bolesnika.

Ključne reči: hiponatrijemija, osteoporoza, natrijum, vitamin D, stare osobe

of calcium homeostasis by enhancing and regulating physiologic calcium absorption by the gut (12), as well as by exerting its effects directly on bone cells. Most of the bone cell types are capable of metabolising 25-OHD to 1,25(OH)₂D₃ (i.e., the active form of the hormone), to stimulate their own biological functions, such as cell proliferation, maturation, mineralisation and bone resorption (12). It is generally acknowledged that an adequate vitamin D status is important to minimize the risk of falling in elderly people, and positive effects on falls and muscle function have been shown in community dwelling older individuals (13–15). A recent meta-analysis involving eight trials and 2,426 subjects showed that doses of vitamin D up to 600 IU were ineffective for lowering the risk of falling, whereas higher doses (ranging from 700 to 1,000 IU) reduced such risk by approximately 20% (16).

In the last decades, vitamin D receptors have been demonstrated to be widespread in brain tissue, and the vitamin D active form has shown neuroprotective effects, including the clearance of amyloid plaques, a hallmark of Alzheimer's disease (17). At least two large prospective studies recently indicated that low vitamin D concentrations may increase the risk of cognitive decline (18), thus representing an adjunctive risk factor for falls in elderly and vitamin D deficient subjects.

Due to the aforementioned similarities between hyponatremia and vitamin D deficiency in increasing the risk of falls, fractures and all-cause mortality, we performed a retrospective, double center investigation to establish whether any association exists between hyponatremia and vitamin D deficiency in the general population.

Materials and Methods

We carried out a retrospective search in the laboratory information systems of the clinical chemistry laboratory of the Academic Hospital of Verona (Italy) and clinical chemistry laboratory of the Academic Hospital of Parma (Italy), to retrieve the combined results of total vitamin D and sodium obtained in all outpatients referred to the two laboratories for routine health check-up during the entire year 2013. In all subjects, the assessment of 25-OHD (Liaison, competitive chemiluminescence assay; DiaSorin, Saluggia, Italy) and sodium (ion-selective electrode method) was performed within 2 hours after blood sample collection. The quality of results was validated by reg-

ular internal quality control procedures and participation in an external quality assessment scheme.

Results were finally expressed as median and interquartile range (IQR). Hyponatremia was defined as serum sodium level lower than 136 mmol/L (5), whereas vitamin D deficiency was defined as a 25-OHD level of less than 20 ng/mL (50 nmol/L) (10). The association between variables was tested with Spearman's correlation and multivariate regression analysis, in which serum sodium concentrations were entered as dependent variables and age, gender and 25-OHD levels as independent variables. Differences between groups were assessed with Wilcoxon-Mann-Whitney and Pearson's χ^2 test with Yates' correction (for categorical variables), using Analyse-it (Analyse-it Software Ltd, Leeds, UK). The odds ratios (ORs) were calculated using MedCalc Version 12.3.0 (MedCalc Software, Mariakerke, Belgium). The study was performed in accordance with the Declaration of Helsinki, under the terms of relevant local legislation.

Results

Combined results of vitamin D and sodium could be retrieved from the laboratory information systems of the two centers for a total of 5097 outpatients throughout the study period (2054 from the laboratory of Parma and 3043 from the laboratory of Verona; 3859 females and 1238 males; mean age 64 ± 17 years). Significantly lower levels of serum sodium were found in vitamin D deficient subjects (140 versus 141 mmol/L; $p < 0.001$), along with a marginally but significantly higher rate of hyponatremia (6.3% versus 5.1%; $p = 0.036$) (Table I). Accordingly, the Odds Ratio (OR) of hyponatremia in vitamin D deficient subjects was 1.26 (95% CI, 0.99 to 1.61). Interestingly, although the calculated mean percentage difference between groups (i.e., 1.0%) was only marginally higher than the inter-individual biologic variation of serum sodium (i.e., 0.7%), the observed

Table I Values of serum sodium stratified according to vitamin D status in a large population of unselected outpatients.

Vitamin D			
	<50 nmol/L	50 nmol/L	p
n	1853	3244	
Age (years)	65 ± 17	64 ± 17	0.053
Gender (F/M)	1363/1853	2496/3244	0.007
Sodium			
Values (mmol/L)	140 (139–141)	141 (139–142)	<0.001
Hyponatremia	117/1853 (6.3%)	165/3244 (5.1%)	0.036

Table II Values of total vitamin D stratified according to serum sodium in a large population of unselected outpatients.

	Sodium		
	<136 mmol/L	136 mmol/L	p
n	282	4815	
Age (years)	66 ± 19	64 ± 17	0.068
Gender (F/M)	213/282	3646/4815	0.995
Vitamin D			
Values (nmol/L)	55 (33–75)	60 (39–82)	0.015
Deficiency	118/282 (41.8%)	1738/4815 (36.1%)	0.030

difference (i.e., 1 mmol/L) was much higher than the minimal clinically important difference for serum sodium calculated in both populations (i.e., 0.06 and 0.04 mmol/L in vitamin D deficient subjects and controls, respectively) (19).

Significantly lower levels of serum 25-OHD were also found in hyponatremic subjects (55 versus 60 nmol/L; $p = 0.015$), along with a marginally but significantly higher rate of vitamin D deficiency (41.8% versus 36.1%; $p = 0.030$) (Table II). Accordingly, the OR of vitamin D deficiency in hyponatremic subjects was 1.27 (95% CI, 1.00 to 1.63). In our region (i.e., the northern part of Italy) it is not usual to consume vitamin D2, so analytic underestimations of 25-OHD due to the characteristics of the Liaison competitive chemiluminescence assay are highly unlikely.

In the univariate regression analysis, a highly significant association was found between the values of sodium and 25-OHD in the entire study population (Spearman's correlation coefficient, 0.11; $p < 0.001$). The association remained statistically significant in the multivariate regression analysis after adjustment for age and gender (beta coefficient, 0.06; $p < 0.001$), as well as in females (Spearman's correlation coefficient, 0.11; $p < 0.001$) and males (Spearman's correlation coefficient, 0.10; $p < 0.001$).

Discussion

The results of this study demonstrate for the first time the existence of a significant positive correlation between the serum levels of sodium and 25-OHD, along with a significant and independent correlation between hyponatremia and vitamin D deficiency in a general population of unselected outpatients, thus opening an intriguing perspective about the potentially meaningful interplay between vitamin D and sodium in human disease. Although some evidence exists that low levels of 25-OHD are weakly associated with

the risk of developing hypertension within 4 and 14 years (19), randomized trials have so far failed to confirm that vitamin D supplementation is effective to lower blood pressure (20).

Plasma renin activity (PRA) and salt sensitivity (SS) of blood pressure (BP) have both been linked to 25-OHD concentrations, and it was also demonstrated that the flux of calcium into vascular smooth muscle cells may be facilitated by the active forms of vitamin D (20). Vitamin D also acts as a proximal inhibitor of the Renin-Angiotensin System (RAS), and higher 1,25(OH)₂D concentrations have been associated with lower plasma renin activity in hypertensive patients (20). Interestingly, an inverse correlation between both 25-OHD and 1,25(OH)₂D and plasma renin and angiotensin II concentrations has been recently reported in a selected cohort of patients referred for coronary angiography (21). Finally, dietary salt loading has been associated with an increase in circulating 1,25OHD concentrations, wherein subjects with the greatest salt-induced increases in 1,25OHD levels were those exhibiting the greatest BP response to salt (20). Conversely, there is some evidence that calcium supplementation can, at least in part, contribute to a good response to antihypertensive drugs in the elderly (22).

Along with the previous evidence, vitamin D is known to be an important modulator of the hemostatic system (23), and low levels have been associated with a higher prevalence of arterial and cardiovascular disease (11), which both represent additional and well established risk factors for syncope and falls.

Nevertheless, it is still uncertain whether hyponatremia and vitamin D deficiency should be considered real pathogenetic factors of falls and fractures or rather simple epiphenomena of these conditions. Further studies are hence needed to elucidate the pathophysiological link between the two parameters,

along with the clinical significance of this potential association (i.e., increased risk of falls, fractures and all-cause mortality with the association of the two metabolic abnormalities). Regardless of the causal or casual nature of this biological interplay, the concentration of serum sodium and 25-OHD should be regarded as a part of a complex and multifactorial impairment that increases the risk of gait disturbances and osteoporosis. Accordingly, it seems reasonable to suggest that serum sodium and 25-OHD levels should be assessed and eventually corrected when the value falls below the lower limit of the reference range.

Unfortunately, no studies have evaluated bone density or incidence of falls and fractures before and after correction of hyponatremia, nor have any prospective investigations aiming to demonstrate potential improvements of clinical outcomes after correction of hyponatremia been published so far. Nevertheless, since sodium is a simple, inexpensive and rapid biochemical test, clinicians should investigate the possible presence of hyponatremia in all elderly patients undergoing therapies with drugs potentially involved in the pathogenesis of hyponatremia, in particular diuretics, psychotropic drugs, and anticonvulsants. The patients with mild chronic hyponatremia should also probably be examined for bone mineral density (BMD) and for falls risk using the available predictive models.

If our observation should be prospectively confirmed, the association of hyponatremia and hypovitaminosis D could be identified and characterized as a new frailty index for elderly patients.

Conflict of interest statement

The authors stated that they have no conflicts of interest regarding the publication of this article.

References

1. Miller M. Hyponatremia and arginine vasopressin dysregulation: mechanisms, clinical consequences, and management. *J Am Geriatr Soc* 2006; 54: 345–53.
2. Holick MF, Siris ES, Binkley N, et al. Prevalence of vitamin D inadequacy among postmenopausal North American women receiving osteoporosis therapy. *J Clin Endocrinol Metab* 2005; 90: 3215–24.
3. Upadhyay A, Jaber BL, Madias NE. Incidence and prevalence of hyponatremia. *Am J Med* 2006; 119: S30–S35.
4. Ellison DH, Berl T. The syndrome of inappropriate antidiuresis. *N Engl J Med* 2007; 356: 2064–72.
5. Adrogué HJ, Madias NE. Hyponatremia. *N Engl J Med* 2000; 342: 1581–9.
6. Hoorn EJ, Rivadeneira F, van Meurs JBJ, et al. Mild Hyponatremia as a Risk Factor for Fractures: The Rotterdam Study. *J Bone Mineral Res* 2011; 26: 1822–8.
7. Verbalis JG, Barsony J, Sugimura Y, et al. Hyponatremia-induced osteoporosis. *J Bone Miner Res* 2010; 25: 54–63.
8. Barsony J, Manigrasso MB, Xu Q, et al. Chronic hyponatremia exacerbates multiple manifestations of senescence in male rats. *Age* 2013; 35: 271–88.
9. Bodyak N, Ayus JC, Achinger S, et al. Activated vitamin D attenuates left ventricular abnormalities induced by dietary sodium in Dahl salt-sensitive animals. *Proc Natl Acad Sci USA* 2007; 104: 16810–5.
10. Dawson-Hughes B, Mithal A, Bonjour JP, et al. IOF po-

- sition statement: vitamin D recommendations for older adults. *Osteoporos Int* 2010; 21: 1151–4.
11. van Schoor NM, Lips P. Worldwide vitamin D status. *Best Pract Res Clin Endocrinol Metab* 2011; 25: 671–80.
 12. Christodoulou S, Goula T, Ververidis A, Drosos G. Vitamin D and bone disease. *Biomed Res Int* 2013: 396541.
 13. Pfeifer M, Begerow B, Minne HW, et al. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community dwelling older individuals. *Osteoporos Int* 2008; 16: 16–28.
 14. Holick FM. The D-lightful vitamin D for health. *J Med Biochem* 2013; 32: 1–10.
 15. Škodrić-Trifunović V, Blanka A, Stjepanović M, Ignjatović S, Mihailović-Vučinić V, Šumarac Z, Buha I, Ilić K. The Health Benefits of Vitamin D Relevant for Tuberculosis. *J Med Biochem* 2014; 33: 301–6.
 16. Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, et al. Fall prevention with supplemental and active forms of vitamin D: A meta-analysis of randomised controlled trials. *BMJ* 2009; 339: b3692.
 17. Dursun E, Gezen-Ak D, Yilmazer S. A novel perspective for Alzheimer's disease: vitamin D receptor suppression by amyloid-beta and preventing the amyloid-beta induced alterations by vitamin D in cortical neurons. *J Alzheimer's Dis* 2011; 23: 207–19.
 18. Llewellyn DJ, Lang IA, Langa KM, Melzer D. Vitamin D and cognitive impairment in the elderly U.S. population. *J Gerontol A Biol Sci Med Sci* 2011; 66: 59–65.
 19. Copay AG, Subach BR, Glassman SD, Polly DW Jr, Schuler TC. Understanding the minimum clinically important difference: a review of concepts and methods. *Spine J* 2007; 7: 541–6.
 20. Vaidya A, Forman JP. Vitamin D and Hypertension. Current Evidence and Future Directions. *Hypertension* 2010; 56: 774–9.
 21. Tomaschitz A, Pilz S, Ritz E, et al. Independent association between 1,25-dihydroxyvitamin D, 25-hydroxyvitamin D and the renin-angiotensin system: the Ludwigshafen Risk and Cardiovascular Health (LURIC) Study. *Clin Chim Acta* 2010; 411: 1354–60.
 22. Cervellin G, Bonino P, Palummeri E, Passeri M. Calcium phosphate and blood pressure: their relationships in a geriatric population. *Am J Nephrol* 1986; 6 Suppl 1: 16–8.
 23. Targher G, Pichiri I, Lippi G. Vitamin D, Thrombosis, and Hemostasis: More than Skin Deep. *Semin Thromb Hemost* 2010; 38: 114–24.

Received: October 23, 2014

Accepted: January 11, 2015